





28-0-10

Fairways Grade DuPont® Acelepryn® Insecticide 0.067% 150SGN

NET WEIGHT 50 LBS. (22.7 KG.) COVERS 14,535 Sq. Ft.

Guaranteed Analysis

Active Ingredients

ACTIVE INGREDIENT:

POUNDS OF NUTRIENT/M SQ. FT.					
Lbs. Per Acre	Lbs. Per 1,000 Sq. Ft.	N	Р	K	
150	3.44	0.96		0.34	
225	5.17	1.45		0.52	
300	6.88	1.93		0.69	

General Information

DuPont® Acelepryn® Insecticide 0.067% 28-0-10 has sufficient residual activity so that applications can be made preceding egg laying activity of the target pests. This product will control white grubs and other listed pest infesting landscape and recreational turfgrass. This product contains 100% of the nitrogen from Polymer-Encapsulated urea (XRT). XRT is a new controlled release technology that delivers a more uniform and predictable release of nutrient for a period of up to 90 days. For Extended Release Technology, trust in XRT to deliver the lasting performance you demand. This product also contains Gro-Gard, a proprietary blend of two bio-active amino acids and co-factors; it activates key amino acids that help regulate nutrient movement between plant cells. And with greater nutrient availability, plants can achieve their full genetic potential for growth and vigor. Gro-Gard protects your turf, optimizes nutrient

uptake and movement within plants, and enhances overall plant health.

The following settings are approximate for applying contents of this package. Settings are based on average walking speed. Reduce settings for slower speed and increase for higher speed. Check your spreader on 250 sq. ft. and adjust to apply ½ of 1,000 sq. ft. rate. PLEASE NOTE: These settings are furnished as a starting guide only.

SPREADER SETTINGS LBS. PER 1,000 SQ. FT.					
SPREADER	3.44 lbs.	5.17 lbs.	6.88 lbs.		
Scott's R8	F-G	H-I	K-L		
Lesco High Wheel	D-E (9)	F-G	H-I (17)		
Spyker 24-22	3-3 ½	4-4 1/2	4 1/2-5		
Earthway Broadcast	14-16	1 <i>7</i> -19	20-22		
Lely	2-4	4-5	5 1/2-6		

Application Instructions

When applying DuPont® Acelepryn® Insecticide 0.067% 28-0-10 you should not apply when the soil is saturated with water because adequate distribution of the active ingredient vertically in the soil profile cannot be achieved under this condition. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

KNOX FERTILIZER COMPANY, INC. P.O. BOX 248 KNOX, IN 46534 (574)772-6275 / FAX (574)772-5878

Shaw's™ & Gro-Gard™ is a registered trademark of Knox Fertilizer Co., Inc. XRT™ is a registered trademark of KMB Technologies, Inc. DuPont® and Acelepryn® are registered trademarks of E.I. DuPont de Nemours & Company